

EVALUATING THE RESPONSE TO FUNGICIDE IN DIFFERENT TILLAGE SYSTEMS

TRIAL OVERVIEW

- Fungicide application to corn is a relatively common practice in Illinois.
- Low commodity prices are calling the return on investment (ROI) for a fungicide application into question.
- Different tillage systems may provide different environments that are more or less preferable to disease development.

RESEARCH OBJECTIVE

- This trial was established to evaluate the yield response to fungicide in different tillage systems.

Location	Soil	Previous Crop	Tillage Type	Planting Date	Harvest Date	Potential Yield/Acre	Planting Rate/Acre
Monmouth, IL	Silt Loam	Corn	Various	04/24/2017	09/29/2017	240 bu/acre	36,000 seeds/acre

SITE NOTES:

- A large land block was divided into three different tillage zones:
 - Vertical Tillage
 - Strip Tillage
 - Conventional Tillage
- Within each of the three tillage zones, two corn products were planted:
 - 108 Day RM SmartStax® RIB Complete® Corn Blend
 - 114 Day RM SmartStax® RIB Complete® Corn Blend
- Each product had treatments consisting of an untreated check and an application of a foliar fungicide that contained strobilurin and triazole active ingredients (A.I.). There were two replications of all treatments. The fungicide was applied at the R1 growth stage.

UNDERSTANDING THE RESULTS




Figure 1. Average yield response (bu/acre) for fungicide application to three tillage systems at Monmouth, Illinois (2017, 2 Replications).




Figure 2. Average yield response (bu/acre) of a fungicide application on two corn products and three tillage systems at Monmouth, Illinois (2017, 2 Replications).



Figure 3. Gray leaf spot, Monmouth, Illinois - 2017.

- Disease incidence was low; however, the disease symptoms that appeared were generally very late in the season and likely had little to no impact on yield.
- No differences in symptomology were seen between the treated and untreated plots.
- Because of these factors, no discernible differences or trends were observed in the final yield results (Figures 1 and 2).
- Three factors are required for disease development: a pathogen, a susceptible host, and favorable environmental conditions.
- In 2017, the cool, dry conditions in July and August likely held disease pressure to a minimum at the Monsanto Learning Center at Monmouth, IL. An example is the minimal number of gray leaf spot lesions found on corn leaves (Figure 3).

WHAT DOES THIS MEAN FOR YOUR FARM?

- Various methods for preventing disease development in corn include planting resistant genetics, crop rotation, and good residue management practices.
- A good scouting program is crucial to identify whether a disease is a problem in any given field.
- If all three factors for disease development are present, a fungicide application may help protect yield potential.

LEGAL STATEMENT

For additional agronomic information, please contact your local brand representative. Developed in partnership with Technology Development & Agronomy by Monsanto.

Monsanto Company is a member of Excellence Through Stewardship® (ETS). Monsanto products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Monsanto's Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for this product. Excellence Through Stewardship® is a registered trademark of Excellence Through Stewardship. **B.t. products** may not yet be registered in all states. Check with your Monsanto representative for the registration status in your state. **IMPORTANT IRM INFORMATION: RIB Complete®** corn blend products do not require the planting of a structured refuge **except** in the Cotton-Growing Area where corn earworm is a significant pest. SmartStax® RIB Complete® corn blend is not allowed to be sold for planting in the Cotton-Growing Area. See the IRM/Grower Guide for additional information. Always read and follow IRM requirements. Individual results may vary, and performance may vary from location to location and from year to year. This result may not be an indicator of results you may obtain as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Roundup Ready technology contains genes that confer tolerance to glyphosate, an active ingredient in Roundup® brand agricultural herbicides. Agricultural herbicides containing glyphosate will kill crops that are not tolerant to glyphosate. RIB Complete®, Roundup Ready®, Roundup® and SmartStax® are trademarks of Monsanto Technology LLC. LibertyLink® and the Water Droplet Design® is a registered trademark of Bayer. Herculex® is a registered trademark of Dow AgroSciences LLC. Respect the Refuge and Corn Design® and Respect the Refuge® are registered trademarks of National Corn Growers Association. All other trademarks are the property of their respective owners.

©2017 Monsanto Company All Rights Reserved.

17101311553 110117LGM

