



DELARO[®]
Complete
 458 SC Fungicide

Corn Gray Leaf Spot Risk Forecast

For period of May 29 – June 26, Date of Issue: May 30, 2021

One-Month Weather Outlook

Over the next 30 days the temperatures will be warmer than normal across the Corn Belt. Precipitation north of I-80 will be near normal, and above normal south of I-80. During the next two weeks temperatures will be above normal with low rainfall north of I-70 and above average precipitation farther south.

Gray Leaf Spot Risk Summary


The risk for gray leaf spot in the Corn Belt is very low in this period of forecast.

Next two-week risk interpretation

Conditions over the next two weeks are unfavorable for the development of gray leaf spot in the Corn Belt as the crop is in early vegetative development. Follow the risk forecasts as corn growth progresses.




ONE-MONTH RISK MAP



WHAT TO DO?	
HIGH	Treat susceptible hybrids and weigh risk factors for other fields
MODERATE	Scout fields and treat as conditions warrant
LOW OR VERY LOW	Monitor risk forecasts

Risk Level



DISCLAIMER
 The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



DELARO[®]
Complete
 458 SC Fungicide

Northern Corn Leaf Blight Risk Forecast

For period of May 29 – June 26, Date of Issue: May 30, 2021

One-Month Weather Outlook

Over the next 30 days the temperatures will be warmer than normal across the Corn Belt. Precipitation north of I-80 will be near normal, and above normal south of I-80. During the next two weeks temperatures will be above normal with low rainfall north of I-70 and above average precipitation farther south.

Northern Leaf Blight Risk Summary


The risk of northern corn leaf blight is low across the Corn Belt at this time.

Next two-week risk interpretation


The risk of NCLB is low in the Corn Belt except for a small area in southwestern Missouri and southeastern Kansas as conditions are generally unfavorable for the disease over the next two weeks. Follow the risk forecasts as the corn crop progresses.



ONE-MONTH RISK MAP



Risk Level



WHAT TO DO?

HIGH	Treat susceptible hybrids and weigh risk factors for other fields
MODERATE	Scout fields and treat as conditions warrant
LOW OR VERY LOW	Monitor risk forecasts

DISCLAIMER
 The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. . **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



Corn Southern Rust Risk Forecast

For period of May 29 – June 26, Date of Issue: May 30, 2021

One-Month Weather Outlook

Over the next 30 days the temperatures will be warmer than normal across the Corn Belt. Precipitation north of I-80 will be near normal, and above normal south of I-80. During the next two weeks temperatures will be above normal with low rainfall north of I-70 and above average precipitation farther south.

Southern Rust Risk Summary

The risk of southern corn rust is very low for the entire Corn Belt at this time.

Next 2-week risk interpretation


Weather conditions are unfavorable to southern corn rust and the disease has yet to move into the northern regions. The risk is very low for the Corn Belt for the next two weeks.



ONE-MONTH RISK MAP



Risk Level



WHAT TO DO?

HIGH	Treat susceptible hybrids and weigh risk factors for other fields
MODERATE	Scout fields and treat as conditions warrant
LOW OR VERY LOW	Monitor risk forecasts

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



DELARO[®]
Complete
 458 SC Fungicide

Corn Common Rust Risk Forecast

For period of May 29 – June 26, Date of Issue: May 30, 2021

One-Month Weather Outlook

Over the next 30 days the temperatures will be warmer than normal across the Corn Belt. Precipitation north of I-80 will be near normal, and above normal south of I-80. During the next two weeks temperatures will be above normal with low rainfall north of I-70 and above average precipitation farther south.

Common Rust Risk Summary


The risk of corn common rust is low to very low in the north except for some areas in Missouri.

Risk interpretation


The risk of corn common rust is low to very low for the next two weeks in the Corn Belt. The disease starts to move into southern Missouri. In areas of yellow conditions are moderately favorable to the disease.



ONE-MONTH RISK MAP



Risk Level



WHAT TO DO?

HIGH	Treat susceptible hybrids and weigh risk factors for other fields
MODERATE	Scout fields and treat as conditions warrant
LOW OR VERY LOW	Monitor risk forecasts

DISCLAIMER
 The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



DELARO[®]
Complete
 458 SC Fungicide

Corn Tar Spot Risk Forecast

For period of May 29 – June 26, Date of Issue: May 30, 2021

One-Month Weather Outlook

Over the next 30 days the temperatures will be warmer than normal across the Corn Belt. Precipitation north of I-80 will be near normal, and above normal south of I-80. During the next two weeks temperatures will be above normal with low rainfall north of I-70 and above average precipitation farther south.

Corn Tar Spot Risk Summary


The risk of corn tar spot in the Corn Belt is low to very low at this time.

Next two-week risk interpretation

Conditions over the next two weeks are unfavorable for corn tar spot development in the Corn Belt as the crop is in early vegetative stages. Follow the risk forecasts as the crop progresses.




ONE-MONTH RISK MAP



WHAT TO DO?	
HIGH	Treat at the first sign of infection in your fields or the disease is developing in neighboring fields
MODERATE	Scout fields and treat as conditions warrant
LOW OR VERY LOW	Monitor risk forecasts

Risk Level



DISCLAIMER
 The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED