



STRATEGO[®]YLD

Soybean Frogeye Leaf Spot Risk Forecast

For period of Aug 21 Sept 19, Date of Issue: August 22, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will be normal east of Mississippi and cooler to the west. Precipitation will be above normal to the west and normal along the east coast. During the next two weeks temperatures are predicted to be higher than normal north of I-20 and normal to the south. Rainfall will be normal east of I-75 and higher to the west.

Frogeye Leaf Spot Risk Summary

The risk of frogeye leaf spot remains moderate or high for much of the South.

Next two-week risk interpretation

During the next two weeks conditions are moderately or highly favorable for the disease. Scout susceptible cultivars in at-risk areas, and if fresh disease lesions are observed on the upper leaves, consider a fungicide treatment if the plants have not reached R5 growth stage.



ONE-MONTH RISK MAP

WHAT TO DO?	
нідн	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Bayer, Bayer Cross and Stratego® YLD are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.







STRATEGO[®]YLD

Asian Soybean Rust Risk Forecast

For period of Aug 21 Sept 19, Date of Issue: August 22, 2021

One-Month Weather Outlook

Over next 30 days temperatures are to be normal east of Mississippi and cooler west. Precipitations are to be above normal west and normal in eastern coast. In next two weeks, temperatures are higher than normal in regions north of I-20 and normal for region south. Precipitations are normal in east of I-75 and higher for region west.

Asian Soybean Rust Risk Summary

The risk of soybean rust is moderate in the Southeast with several high risk areas.

Next two-week risk interpretation

During the next two weeks the weather conditions are moderately favorable for the disease in the Southeast. Scout late-planted soybeans in at-risk areas. Consider treating fields where the disease is observed and the plants have not reached R5 growth stage.



ONE-MONTH RISK MAP

WHAT TO DO?	
нідн	Consider treatment if SBR is confirmed in the area
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Bayer, Bayer Cross and Stratego® YLD are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

