



Pre-Plant Nitrogen Optimization for Corn

Key Points

- Fertilization method, application timing, and fertilizer cost should be evaluated to minimize nitrogen (N) loss while meeting crop needs.
- All N fertilizer formulations have the potential for loss. Understanding these loss mechanisms can help farmers make informed choices for each unique situation.
- Denitrification, leaching, and volatilization are three processes that can lead to N loss from a pre-plant application.

Potential Nitrogen Loss

Denitrification

Denitrification is the process by which soil bacteria that thrive in water-saturated (anaerobic) soils convert nitrate into gaseous forms of N. This bacterial conversion of nitrate into nitrogen gas can result in as much as 5% nitrate loss per day. Losses will be higher with warmer soil temperatures.

Leaching

Leaching occurs when nitrate moves downward in the soil profile, out of the root zone, due to excessive rainwater or irrigation. Fertilizers that contain urea ammonium nitrate (UAN) are highly susceptible to leaching, especially when applied to coarse-textured soils.

Volatilization

Volatilization can occur when urea-based fertilizers are surface-applied and not incorporated. Urease enzymes in the soil and plant residue convert urea to free ammonia gas. Up to 20% of urea-based N can volatilize within a week of application, especially on warm sunny days with moist soil surface conditions.

Volatilization can be inhibited or delayed through the use of products such as nitrification inhibitors, urease inhibitors, and polymer-coated urea.



(Figure 1). Anhydrous ammonia applications in the spring.

Pre-Plant Nitrogen Optimization for Corn

Nitrogen Application Timing

Fall applications have the greatest risk of N loss. Anhydrous ammonia is recommended for situations in which fall N applications are preferred, because it has the lowest loss risk of any N fertilizer. Anhydrous ammonia should be applied with a nitrification inhibitor when soil temperature is less than 50 °F (10 °C). A nitrification inhibitor works by slowing down the conversion of ammonium to nitrate in the soil, reducing the potential risk of N loss.

Spring pre-plant applications are generally preferable due to their lower potential for N loss and improved timing relative to rapid plant N uptake. However, spring can also be a time of large potential N loss due to wet soils. If N is applied more than two weeks prior to planting, anhydrous ammonia is recommended to help reduce the risk of loss. Nitrogen loss potential due to leaching is highest when applied to sandy or coarse-textured soils.

Strip till pre-plant fertilizer applications can apply both a N source and a second fertilizer, or combination of fertilizers, six to eight inches deep. This can be done in the fall or in the spring prior to planting. This one-trip tillage operation tills a narrow strip to prepare a seed bed while positioning pre-plant fertilizer several inches below the seed slice where the seed will be placed at planting. Placing a band of fertilizer into the future root zone increases plant utilization, especially if the soil has very low nutrient availability.

At-planting or in-season sidedress applications may be required if soil conditions or workload prevent a pre-plant application of N fertilizer, pushing the N application into the at-planting or in-season sidedress window. If such an application is required, note that any N source can be susceptible to denitrification if the soil is saturated after the application and after the N has been converted to nitrate by soil organisms. If available, fertigation using an overhead sprinkler irrigation can also be effective.

Caution – Avoid placing high rates of N or potassium fertilizer in direct contact with seed to prevent potential seedling injury from salt burn.

Nitrogen Sources and Application

Ammonium and nitrate are the two forms of N used by plants. Other forms of N must be converted by soil organisms to one of these forms before plants can use them, and so the N source should be considered when determining which application method to use.

Anhydrous ammonia is the densest nitrogen source. All other nitrogen fertilizer sources are made from anhydrous ammonia, with additional manufacturing processes that reduce the concentration and typically increase the cost per unit of N (though cost can be affected by seasonal supply and demand). However, all other N sources are generally easier to store and apply, and are safer to handle than anhydrous ammonia.

Table 1. Nitrogen (N) source and application.

N-Source Analysis	Common Name	N Form	Time of Application	Advantage of Product	Primary Method of N Loss	Use with N Stabilizer/Inhibitor
82-0-0	Anhydrous Ammonia	Gas/Liquid	Fall, Pre-plant, Sidedress	Nitrogen Dense	Volatilization	Yes
46-0-0	Urea	Dry Pellet	Pre-Plant, Band	Good Storage and Handling	Volatilization/Leaching	Yes
28/32-0-0	Urea Ammonium Nitrate (UAN)	Liquid	Pre-plant, Sidedress	Relatively Easy to Handle and Apply	Volatilization/Leaching	Yes



Pre-Plant Nitrogen Optimization for Corn

Determining the Nitrogen Application Rate

The rate of N applied is an important variable, because of the related economic and environmental issues. Determining the proper N application rate can be difficult due to the effects of temperature and precipitation on the release of N (mineralization) from the soil over time and after an application, and because microbial activity releases more N under good corn-growing conditions.

Before determining the amount of N to apply, the producer must:

- Estimate the amount of N already in the soil that will be available for corn crop use
- Determine the total N needed for the expected crop yield
 - » As yield goals increase, the amount of N required will also increase



Figure 2. A growing corn crop that is nitrogen deficient.

Crop nitrogen needs and the corresponding N applications must be calculated every year and for every field. A common calculation to determine the total N needed for an acre of corn is multiplying the yield goal for the projected crop year by a factor of 1.0 to 1.2 pounds of N per bushel per acre, and applying N based upon yield goals and soil test levels has been a common method used in western corn growing areas.

Alternatively, several universities in the Corn Belt recommend using a “maximum return to N” (MRTN) formula to determine an appropriate N application rate. These calculations are made for a single N cost and corn price combination; different price combinations can be placed on the same graph to compare additional N cost and corn price combinations. Different states and different crop rotations are used to formulate different MRTN graphs for those situations. For more information on MRTN calculations, see the Corn Nitrogen Rate Calculator at <https://www.cornnratecalc.org/> or the Ontario Corn Nitrogen Calculator at <http://www.gocorn.net/>.

Consider all N sources when making application decisions, including any previous legume crops and manure applications, and credit the N supplied from those sources to reduce the amount of nitrogen fertilizer needed. Using variable rate technology to apply more N in fields or areas of fields where there is greater potential for response can also help increase yield potential.

Pre-Plant Nitrogen Optimization for Corn

Additional Resources

Scharf, P. and Lory, J. 2018. Best management practices for nitrogen fertilizer in Missouri. University of Missouri Extension. <https://extension.missouri.edu/publications/ipm1027>

Nowatzki, J., Endres, G., DeJong-Hughes, J., and Aakre, D. 2017. Strip till for field crop production. North Dakota State University. AE1370.

<https://www.ag.ndsu.edu/publications/crops/strip-till-for-field-crop-production>

Sawyer, J. 2021. Corn nitrogen rate calculator. Iowa State University Agronomy and Extension Outreach.

<https://crops.extension.iastate.edu/corn-nitrogen-rate-calculator>

Legal Statements

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.

Performance may vary, from location to location and from year to year, as local growing, soil and environmental conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on their growing environment.

The recommendations in this material are based upon trial observations and feedback received from a limited number of growers and growing environments. These recommendations should be considered as one reference point and should not be substituted for the professional opinion of agronomists, entomologists or other relevant experts evaluating specific conditions.

Bayer and Bayer Cross are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. ©2024 Bayer Group. All rights reserved. 1213_58761

